

THE CLUSTER-MERGER SHOCK IN 1E 0657–56: FASTER THAN THE SPEEDING BULLET?

MILOŠ MILOSAVLJEVIĆ¹, JUN KODA¹, DAISUKE NAGAI², EHUD NAKAR², AND PAUL R. SHAPIRO¹

Draft version February 5, 2008

ABSTRACT

Shock waves driven in the intergalactic medium during the merging of galaxy clusters have been observed in X-ray imaging and spectroscopy. Fluid motions inferred from the shock strength and morphology can be compared to the cold dark matter (CDM) distribution inferred from gravitational lensing. A detailed reconstruction of the CDM kinematics, however, must take into account the nontrivial response of the fluid intracluster medium to the collisionless CDM motions. We have carried out two-dimensional simulations of gas dynamics in cluster collisions. We analyze the relative motion of the clusters, the bow shock wave, and the contact discontinuity and relate these to X-ray data. We focus on the “bullet cluster,” 1E 0657–56, a near head-on collision of unequal-mass clusters, for which the gas density and temperature jumps across the prominent bow shock imply a high shock velocity $4,700 \text{ km s}^{-1}$. The velocity of the fluid shock has been widely interpreted as the relative velocity of the CDM components. This need not be the case, however. An illustrative simulation finds that the present relative velocity of the CDM halos is $\sim 16\%$ lower than that of the shock. While this conclusion is sensitive to the detailed initial mass and gas density profile of the colliding clusters, such a decrease of the inferred halo relative velocity would increase the likelihood of finding 1E 0657–56 in a Λ CDM universe.

Subject headings: dark matter — galaxies: clusters: general — galaxies: clusters: individual (1E 0657–56) — intergalactic medium — shock waves — X-rays: galaxies: clusters

1. INTRODUCTION

During the growth of structure in the universe, nonlinear cold dark matter (CDM) halos collide and merge hierarchically. When halos collide, the dark matter of one can, if collisionless, pass through that of the other, unlike the fluid baryonic gas in the halos. Since the collisions are frequently supersonic for this gas, it passes through shocks where it heats and virializes in the final halo. Currently, an X-ray surface brightness and temperature discontinuity identifiable as a merger shock with Mach number significantly above unity has been detected with *Chandra* in two galaxy clusters: 1E 0657–56 (Markevitch et al. 2002; Markevitch 2006) and A520 (Markevitch et al. 2005). In both cases, X-ray maps exhibit a bow-shock like temperature and density jump. For 1E 0657–56, the spatial segregation of the X-ray emitting plasma from the peaks of the mass distribution detected with gravitational lensing has been interpreted as the first direct proof of the existence of dark matter (Clowe et al. 2006).

The strength and geometry of these shock fronts, and the structure of the contact discontinuity, depend on the kinetic energy and the detailed gravitational and gas density profiles of the merging components. If one could recover the collisionless CDM dynamics from the fluid dynamics of the X-ray emitting gas, then the kinematics of merger shocks would be a powerful probe of the nature and clustering of CDM (e.g., Markevitch et al. 2005; Clowe et al. 2006). However, the two components are coupled only by gravity and behave differently in the merging process. Recently, it has been assumed that the gas shock velocity inferred from the X-ray measurements of 1E 0657–56 equals that of the colliding clusters, implying a rare merger event that might be in conflict with the statistical expectations of the CDM model. We critically examine this assumption by simulating gas dynamics in clus-

ter mergers and studying the response of the intergalactic medium (IGM).

In the cluster-merger in 1E 0657–56¹, transverse motion of the merger shock ($\approx 4,700 \text{ km s}^{-1}$; Markevitch 2006) greatly exceeds relative radial motion of galaxies in the two components ($\approx 600 \text{ km s}^{-1}$; Barrera et al. 2002). This suggests that the shock was driven in the IGM of the larger component by a near head-on supersonic passage of the smaller component. Inside the bow shock, another bow-shaped discontinuity with a reverse temperature jump (lower temperature on the convex side) is seen; this has been interpreted as the contact discontinuity separating the shocked and ram-pressure stripped IGM of the two merging components. Takizawa (2005, 2006) recovered the three basic features (a bow shock, a contact discontinuity, and ram-pressure stripping) in hydrodynamical simulations of cluster collisions, but did not address the relative kinematics of the CDM and the fluid component.

We study the formation and propagation of the merger shock wave and the evolution of the contact discontinuity, and relate these observable features of X-ray surface brightness and temperature maps to CDM kinematics. In § 2, we describe the numerical method, report the general features of the simulations, and analyze a particular run that approximates the observed 1E 0657–56 (the conclusions also apply to other cluster mergers). In § 3, we discuss implications for CDM clustering. The standard cosmological parameters consistent with WMAP (Spergel et al. 2006) are assumed throughout.

2. SIMULATIONS OF CLUSTER MERGER SHOCKS

2.1. The Algorithm and Initial Conditions

The simulations were carried out with the thoroughly-tested Eulerian code ASC FLASH (Fryxell et al. 2000) in two spatial dimensions. Its adaptive mesh refinement capability al-

¹ Department of Astronomy, University of Texas, 1 University Station C1400, Austin, TX 78712.

² Theoretical Astrophysics, Mail Code 130-33, California Institute of Technology, 1200 East California Boulevard, Pasadena, CA 91125.

¹ With an average X-ray temperature of $14.1 \pm 0.2 \text{ keV}$ (Markevitch et al. 2002; Markevitch 2006; Andersson et al. 2006; Markevitch & Vikhlinin 2007), the cluster 1E 0657–56 at redshift $z = 0.296$ (Tucker et al. 1998) is the hottest and X-ray brightest known cluster.

lowed us to simulate a large spatial domain with dimensions of 20 Mpc and simultaneously achieve a spatial resolution of ~ 1 kpc at fluid discontinuities. The simulations were carried out in cylindrical (r, z) coordinates in an inertial frame. Axial symmetry restricts the simulations to head-on cluster collisions. To avoid artifacts which can result when a perfectly head-on collision occurs with cuspy halo profiles and to approximate the conditions in non-head-on collisions, the central dark matter and gas density of the larger cluster was artificially reduced. The gravitational potential of each of the clusters is spherically symmetric and a fixed function of distance from the center. The clusters are placed at separation D_{12} and are allowed to move under each other's gravity. The acceleration of each cluster is evaluated at its center. The effect of the Hubble flow on cluster motion is ignored because it only affects the early infall, while we are interested in the dynamics of the two halos and their gas content at pericenter passage. We also ignore the dynamical friction drag that reduces the halo velocity during pericenter passage. The fluid flow and its relation to CDM dynamics within a few hundred kiloparsecs from the center of the smaller halo should not be different in the presence of dynamical friction, because the gravitational field of the dynamical wake that trails the smaller CDM halo will not have strong gradients (tides) on such scales. Finally, we ignore the tidal stripping of the smaller CDM halo which could be substantial on larger scales than are relevant here.

For the mass distribution associated with the gravitational potential we consider the Navarro et al. (1997) profile (“NFW”) $\rho_{\text{NFW}} = \rho_0(r/r_s)^{-1}(1+r/r_s)^{-2}$, and the “core” profile $\rho_{\text{core}} = \rho_0(1+r/r_s)^{-3}$. The profiles were normalized such that the mass enclosed within a fixed radius r_{500} equals the mass of the cluster M_{500} . The gas density profile was allowed to differ from that of the CDM profile. At the start of each simulation, monatomic gas with adiabatic polytropic equation of state ($\gamma = 5/3$) is set in hydrostatic equilibrium within each halo. Explicit viscosity, heat conduction, cooling, and self-gravity of the gas are ignored. Runs were carried out for a variety of merger velocities and density profiles that included NFW and core CDM profiles, as well as cuspy (divergent) and non-cuspy gas density profiles. In divergent CDM density profiles, a small constant-density core (20 kpc) in hydrostatic equilibrium was applied in the center to resolve the fluid gradients with multiple resolution elements. The cluster gas fraction was $f_{\text{gas}}\Omega_b/\Omega_m$, where $\Omega_b/\Omega_m \approx 0.16$ is the cosmological baryon abundance and $f_{\text{gas}} \approx 0.7$ (Afshordi et al. 2006).² The mean molecular weight was $\mu = 0.59$, corresponding to 75% hydrogen and 25% helium by mass, fully ionized. Numerical convergence was ascertained by resimulation at increased spatial resolution.

2.2. General Features of Cluster Merger Shocks

Two prominent features seen in all simulations are a bow shock wave, and a contact discontinuity (cold front). The formation of a shock wave during cluster merger was discussed by Markevitch et al. (2000); our results are consistent with their picture. Opening angle and the radius of curvature of the nose of the bow shock are sensitive to the details of the simulation, but both are larger than those expected in steady-state bow shocks driven by spherical hard spheres moving with a constant velocity in a uniform medium (e.g., Farris & Russell




FIG. 1.— Temperature in the meridional plane after cluster pericenter passage. The larger CDM halo is located at $(r, z) = (0, 0)$. The circles indicate the scale radii r_s of the two halos.

1994; Vikhlinin et al. 2001, and references therein). Material on the convex side of the contact discontinuity originates in the smaller cluster. During the collision, a reverse shock propagates into the smaller cluster; a fraction of the smaller cluster traversed by the reverse shock is heated, but the dense core of the cluster remains cold as the reverse shock weakens inside it. The wings of the contact discontinuity are Kelvin-Helmholtz (KH) unstable (in real clusters, the KH instability will be modified by magnetic stresses). A nonlinear instability disrupts the coherence of the surface of the discontinuity and creates a narrowing in the axial diameter of the discontinuity (a neck) just behind the nose, giving it a mushroom-like or bullet-like appearance, as evident in the temperature map in Figure 1. At later times (not shown), the Rayleigh-Taylor instability eats into and disrupts the bullet from the rear, as predicted by Markevitch & Vikhlinin (2007).

In most simulations, during pericenter passage, the shock and contact discontinuity are slower than the relative velocity of the two CDM halos (shock velocity is calculated in the local rest frame on the upstream side of the shock). This is due to the ram-pressure force acting on the gas in the center of the smaller cluster but not on the halo. In many runs some degree of ram-pressure stripping is evident in the center of the smaller cluster (we call the smaller cluster ram-pressure stripped when cold gas originating in the smaller cluster has been expelled from the center of the smaller halo). However, later, as the halos climb out of each other's gravitational potential well and decelerate, but the shock and the contact discontinuity do not decelerate appreciably over a longer period (Fig. 2a). The lack of deceleration of the shock and the cold front was noticed, though not quantified, in previous simulations of cluster mergers (Markevitch & Vikhlinin 2007, and references therein). It could result from the drop in ram pressure as the cold bullet propagates into a thinning cluster at

² All gas-dynamical results are independent of this assumed value of f_{gas} , except that the X-ray surface brightness $\propto f_{\text{gas}}^2$.




FIG. 2.— (a) Velocity of the shock (solid) and of the contact discontinuity (dashed) relative to the pre-shock gas upstream, and relative velocity of the CDM halos (dot-dashed), all as functions of the time-varying separation between the halos $D_{12} = z_2 - z_1$. (b) The density (solid) and temperature jump (dashed) at the shock, and the shock Mach number (dot-dashed). (c) Distance of the shock (solid) and the contact discontinuity (dashed) from the center of the smaller halo.

mosphere and from the gravitational tide of the halo. After the cold gas originally in the smaller cluster has been ram-pressure stripped, the stripped gas lags behind the smaller halo, but eventually its velocity becomes larger than that of the halo, which implies that the gas can later catch up with the halo again.

Before proceeding to discuss a specific simulation approximating the observed bullet cluster, we strongly emphasize that the shock kinematics and morphology are *extraordinarily sensitive* to the parameters of the simulation. In particular, a small variation ($\sim 25\%$) in the cluster mass ratio or density profile can lead to vastly different degrees of ram-pressure stripping, or the opening angle of the bow shock.

2.3. A Model of 1E 0657–56

General results discussed in § 2.2 are illustrated by a specific run that came the closest to reproducing the observed properties of 1E 0657–56. Cluster masses in the run were $M_{500,1} = 1.27 \times 10^{15} M_\odot$ and $M_{500,2} = 2.54 \times 10^{14} M_\odot$, cluster radii were $r_{500,1} = 1.5$ Mpc and $M_{500,2} = 1$ Mpc, and the cluster scale radii were $r_{s,1} = 500$ kpc and $r_{s,2} = 333$ kpc.³ The density

³ The mass ratio of the two halos in the simulation, $M_{500,1}/M_{500,2} = 5$, is about a half of the ratio ~ 10 inferred from lensing data (Clowe et al. 2004, 2006; Bradač et al. 2006). Some fraction of the smaller cluster’s initial mass will have been tidally stripped during the merger, possibly reconciling our initial conditions with lensing. The total luminosity L_{tot} of galaxies in each cluster can be used to estimate the cluster mass via the relation $L_{\text{tot}} \propto M_{500}^\beta$ (Lin et al. 2004; Miller et al. 2005; Cooray & Milosavljević 2005), where $\beta \lesssim 1$ is a bandpass-dependent factor. For the R -band we adopt $\beta_R = 0.65$. Barrena et al. (2002) report $L_{\text{tot},1,R} = 10^{12} L_\odot$ and $L_{\text{tot},2,R} = 0.2 \times 10^{12} L_\odot$, implying a mass ratio of $M_{500,1}/M_{500,2} \approx 12$, in agreement with lensing. The difference of the mass ratios may also be attributed to a slightly non-head-on collision in 1E 0657–56. Indeed, the lensing maps are consistent with a northerly passage of the smaller cluster inducing the observed SW–NE tidal




FIG. 3.— Temperature profile on the central axis (left panel, solid line); emissivity-weighted integrated temperature profile (left panel, dotted line); and 0.8–4 keV surface brightness on the central axis (right panel). The center of the larger CDM halo is located at $z = 0$. The bow shock at $z \sim 0.94$ Mpc is split into a main shock and a weaker subshock.

profile of the larger cluster was the non-divergent ρ_{core} with central density $\approx 2.3 \times 10^{-3} M_\odot \text{pc}^{-3}$ (again, to approximate the conditions in a non-head-on merger with pericenter passage at a distance $\sim r_{s,1}$), and, in the smaller cluster, it was the NFW profile ρ_{NFW} . The gas density in both clusters was proportional to the CDM density, $\rho_{\text{gas}} = (1 + \Omega_m/f_{\text{gas}}\Omega_b)^{-1} \rho_{\text{CDM}}$. The halos initiated infall from relative rest at separation $D_{12} = 4.6$ Mpc and their maximum relative velocity at pericenter was $V_{\text{max}} \approx 5,270 \text{ km s}^{-1}$.⁴

Figure 1 shows a snapshot of the meridional temperature distribution when the distance of the shock to the larger CDM halo is about 1 Mpc. The larger and smaller cluster are moving vertically upward and downward in the figure, respectively. The cold gas bullet is ~ 100 kpc in radius, excluding visible fingers of cold gas embedded in the hot post-shock medium. The center of the smaller halo is located ahead (below) the contact discontinuity, and almost exactly at the location of the bow shock, as found in X-ray and lensing maps of 1E 0657–56 (Clowe et al. 2004, 2006; Bradač et al. 2006). This is also evident in Figure 2c showing the position of the bow shock and the contact discontinuity relative to the position of the smaller halo. Ram-pressure stripping at the center of the smaller cluster occurs at $D_{12} = 0$, but the bow shock and the contact discontinuity catch up with the halo at $D_{12} \approx 1.2$ Mpc and ≈ 1.5 Mpc respectively.

The pre- and post-shock axial temperatures are 12 keV and 34 keV, respectively (see Figure 3). Figure 2b shows the evolution of the density and temperature jumps across the shock, and its Mach number, as a function of the halo separation. The Mach number is ≈ 3 when the separation equals the projected separation of lensing density peaks $D_{12} \approx 720 \pm 25$ kpc (Clowe et al. 2004, 2006; Bradač et al. 2006). The temperatures and shock strength are compatible with the results of Markevitch (2006), who, for a $\gamma = 5/3$ polytropic equation of state and instant electron-ion equilibration, derived the Mach number from deprojected density and temperature jumps in the bow shock in 1E 0657–56. The separation of the shock and the contact discontinuity is ~ 70 kpc, somewhat smaller than in the observed system. Figure 4 shows the predicted X-ray surface brightness map for the simulated cluster.

Figure 2a shows the evolution of the velocities of the shock, the contact discontinuity, and the relative velocity of the two CDM halos. While the cluster velocity decreases as the small

distortion in the larger cluster. Then, the lower density column met by the small cluster would result in equivalent shock dynamics if the smaller cluster were smaller than in the simulation.

⁴ All dimensionless numbers (such as the shock Mach number, and the gas-to-dark matter velocity ratio) and distances (such as the radius of curvature of the shock) are invariant under a uniform scaling of the CDM density that preserves lengths. The scaling preserves the mass to temperature ratio.




FIG. 4.— Surface brightness map of the simulated bullet cluster in the 0.8–4 keV band, assuming a frequency-independent effective area of 400 cm².

halo climbs the potential of the larger halo, the shock velocity remains constant. The lack of deceleration of the shock can be explained by a combination of factors; in addition to gravitational forces, the negative density and temperature gradients into which the shock propagates tend to increase its strength. At $D_{12} = 720$ kpc, the velocity of the shock is $V_{sh} \approx 4,800$ km s⁻¹, consistent with $V_{sh} \sim 4,740^{+710}_{-550}$ km s⁻¹ inferred for 1E 0657–56 by Markevitch (as cited in Farrar & Rosen 2006, private communication). However, the relative velocity of the halos in our simulation is much less, $V_{sub} \sim 4,050$ km s⁻¹.⁵

3. DISCUSSION AND CONCLUSIONS

Recent analyses of 1E 0657–56 (e.g., Hayashi & White 2006; Clowe et al. 2006; Bradač et al. 2006; Farrar & Rosen 2006) assume that the shock velocity equals the instantaneous relative velocity of the CDM halos, and notice that it is much larger than the virial velocity of the larger cluster. Hayashi & White (2006, hereafter HW06) and Farrar & Rosen (2006, hereafter FR06) estimate the probability that a halo in the Λ CDM universe has a subhalo with velocity larger than that of the bow shock in 1E 0657–56. We assess a correction that should be applied to the inferred probabilities to account for the relative motion of the shock and the subhalo. HW06 calculate the likelihood of finding a halo-subhalo pair with such relative velocity by extrapolating results of the Millenium cosmological N -body

⁵ If the true halo separation is larger than the projected separation 720 kpc, the predicted subcluster velocity V_{sub} will be $< 4,050$ km s⁻¹.

⁶ We do not test the accuracy of the cited values of V_{200} , but the lensing data seem to indicate that HW06’s value of V_{200} is an overestimate.

simulation. They find that it decreases rapidly with V_{sub} , $\log[N_1(> V_{sub})/N_{hosts}] = -(V_{sub}/v_{10}V_{200})^\alpha$, where $v_{10} \approx 1.55$ and $\alpha \approx 3.3$. The probability depends on the assumed values of V_{sub} and V_{200} , in which HW06 and FR06 differ.⁶ HW06 adopt a shock velocity of $V_{sh} \approx 4,500^{+1100}_{-800}$ km s⁻¹ (Markevitch et al. 2004) as the relative velocity V_{sub} of the two halos and adopt $V_{200} \approx 2,380$ km s⁻¹ for the virial velocity. With these, they infer that the probability of most massive subhalo having such a velocity is $N_1(> V_{sub}) \approx 1\%$, whereas when they adopt a lower shock velocity consistent with uncertainties of $V_{sub} = 3,700$ km s⁻¹, the probability is $N_1(> V_{sub}) \approx 10\%$. In our simulation (§ 2.3), we find that, when the halos are separated by the observed projected distance $D_{12} = 720$ kpc, the velocity of the shock and that of the subhalo differ by $(V_{sh} - V_{sub})/V_{sub} \approx 16\%$. Thus, the probabilities reported by HW06 should be corrected and should read $\approx 8\%$ and $\approx 27\%$, respectively. FR06 adopt a somewhat higher shock velocity $V_{sh} \approx 4,740$ km s⁻¹ and significantly lower virial velocity $V_{200} \approx 1,740$ km s⁻¹. These values imply a very low probability of $\approx 3.7 \times 10^{-7}$.⁷ The probability is significantly higher but remains very low, $\approx 2.4 \times 10^{-4}$, when the 16% correction for the difference between the shock velocity and CDM halo velocity is taken into account.

In conclusion, our 2D hydrodynamical simulations of merging galaxy clusters show that, in a case like the “bullet cluster” 1E 0657–56, the halo collision velocity need not be the same as the intergalactic gas shock velocity. While the kinematics of the shock is sensitive to the details of the cluster structure, the instantaneous shock velocity can exceed the relative velocity of CDM halos by at least $\sim 16\%$. Any attempt to relate shock kinematics to the details of hierarchical clustering and halo assembly must take into account the nontrivial response of the IGM. Published estimates of the likelihood of finding a configuration resembling 1E 0657–56 in a Λ CDM universe may require upward revision by a factor of 2 to 700.

The research was supported in part by the Sherman Fairchild Foundation and NASA Astrophysical Theory Program grant NNG04G177G. The software used in this work was developed in part by the DOE-supported ASC / Alliance Center for Astrophysical Thermonuclear Flashes at the University of Chicago.

⁷ HW06 and FR06 also report $\times 5$ lower probabilities after counting only post-collision systems.

REFERENCES

- Afshordi, N., Lin, Y.-T., Nagai, D., & Sanderson, A. J. R. 2006, preprint (astro-ph/0612700)
 Andersson, K. E., Peterson, J. R., & Madejski, G. M. 2006, preprint (astro-ph/0612038)
 Barrena, R., Biviano, A., Ramella, M., Falco, E. E., & Seitz, S. 2002, A&A, 386, 816
 Bradač, M., et al. 2006, ApJ, 652, 937
 Clowe, D., Gonzalez, A., & Markevitch, M. 2004, ApJ, 604, 596
 Clowe, D., Bradač, M., Gonzalez, A. H., Markevitch, M., Randall, S. W., Jones, C., & Zaritsky, D. 2006, ApJ, 648, L109
 Cooray, A., & Milosavljević, M. 2005, ApJ, 627, L85
 Farrar, G. R., & Rosen, R. A. 2006, preprint (astro-ph/0610298)
 Farris, M. H., & Russell, C. T. 1994, J. Geophys. Res., 99, 17681
 Fryxell, B., et al. 2000, ApJS, 131, 273
 Hayashi, E., & White, S. D. M. 2006, MNRAS, 370, L38
 Lin, Y.-T., Mohr, J. J., & Stanford, S. A. 2004, ApJ, 610, 745
 Markevitch, M., et al. 2000, ApJ, 541, 542
 Markevitch, M., Gonzalez, A. H., David, L., Vikhlinin, A., Murray, S., Forman, W., Jones, C., & Tucker, W. 2002, ApJ, 567, L27
 Markevitch, M., Gonzalez, A. H., Clowe, D., Vikhlinin, A., Forman, W., Jones, C., Murray, S., & Tucker, W. 2004, ApJ, 606, 819
 Markevitch, M., Govoni, F., Brunetti, G., & Jerius, D. 2005, ApJ, 627, 733
 Markevitch, M. 2006, ESA SP-604: The X-ray Universe 2005, 723
 Markevitch, M., & Vikhlinin, A. 2007, ArXiv Astrophysics e-prints, arXiv:astro-ph/0701821
 Miller, C. J., et al. 2005, AJ, 130, 968

Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, *ApJ*, 490, 493
Spergel, D. N., et al. 2006, preprint (astro-ph/0603449)
Tucker, W., et al. 1998, *ApJ*, 496, L5
Takizawa, M. 2005, *ApJ*, 629, 791

Takizawa, M. 2006, *PASJ*, 58, 925
Vikhlinin, A., Markevitch, M., & Murray, S. S. 2001, *ApJ*, 551, 160